



NAUSEA & VOMITING	
ADULT	PEDIATRIC
BLS Procedures	
Assess airway, suction airway as needed. Assess vitals. Obtain SpO2 Oxygen. Titrate to SpO2 to 94% or higher.	
EMT Optional Scope	
<u>Administration of Ondansetron (Zofran) by EMT is only approved for EMT's that have successfully completed a Stanislaus County EMS Agency approved training.</u>	
ADULT PATIENTS ONLY (18 years and up) Ondansetron (Zofran) 4 mg PO <ul style="list-style-type: none"> May repeat once after 15 min 	Pediatric Ondansetron (under 18 yrs. old) not approved for EMT's
First Trimester Pregnancy (less than 12 weeks)	
Withhold Ondansetron (Zofran) in First Trimester Pregnancy unless already using via prescription.	
ALS Standing Orders	
IV access as needed. Obtain ECG.	
Ondansetron (Zofran) 4 mg PO/IM/IV/IO <ul style="list-style-type: none"> Slow IV push over 1 minute. Single max dose 4 mg. May repeat every 10 minutes. Total max dose 12 mg 	<p style="text-align: center;"><u>2-3 Years old</u></p> Ondansetron (Zofran) 2 mg IM/IV/IO <ul style="list-style-type: none"> Slow IV Push over 1 minute. No repeat
<u>First Trimester Pregnancy (less than 12 weeks)</u> Withhold Ondansetron (Zofran) in First Trimester Pregnancy unless already using via prescription.	<p style="text-align: center;"><u>4 Years and older</u></p> Ondansetron (Zofran) 4 mg PO/IM/IV/IO <ul style="list-style-type: none"> Slow IV push over 1 minute. No repeat
Special Considerations	
<ul style="list-style-type: none"> Nausea and Vomiting are signs and symptoms of many other illnesses and or diseases. Have a high suspicion of cardiac related conditions or dysrhythmias. 	
Base Hospital Orders Only	
Contact Base Hospital Physician for Doses that exceed written protocol, or for patients under 2 years old.	